

JAMA | **Original Investigation** | CARING FOR THE CRITICALLY ILL PATIENT

Effect of Postextubation High-Flow Nasal Oxygen With Noninvasive Ventilation vs High-Flow Nasal Oxygen Alone on Reintubation Among Patients at High Risk of Extubation Failure A Randomized Clinical Trial

Arnaud W. Thille, MD, PhD; Grégoire Muller, MD; Arnaud Gacouin, MD; Rémi Coudroy, MD; Maxens Decavèle, MD; Romain Sonnevile, MD, PhD; François Beloncle, MD; Christophe Girault, MD; Laurence Dangers, MD; Alexandre Lautrette, MD, PhD; Séverin Cabasson, MD; Anahita Rouzé, MD; Emmanuel Vivier, MD; Anthony Le Meur, MD; Jean-Damien Ricard, MD, PhD; Keyvan Razazi, MD; Guillaume Barberet, MD; Christine Lebert, MD; Stephan Ehrmann, MD, PhD; Caroline Sabatier, MD; Jeremy Bourenne, MD; Gael Pradel, MD; Pierre Bailly, MD; Nicolas Terzi, MD, PhD; Jean Dellamonica, MD, PhD; Guillaume Lacave, MD; Pierre-Éric Danin, MD; Hodanou Nanadoumgar, MD; Aude Gibelin, MD; Lassane Zanre, MD; Nicolas Deye, MD, PhD; Alexandre Demoule, MD, PhD; Adel Maamar, MD; Mai-Anh Nay, MD; René Robert, MD, PhD; Stéphanie Ragot, PharmD, PhD; Jean-Pierre Frat, MD; for the HIGH-WEAN Study Group and the REVA Research Network

JAMA.2019;322(15):1465-1475.

2019年11月26日

慈恵ICU Journal Club

初期研修医2年目 石崎 俊太